arXiv:1201.4222v2 [nlin.SI] 31 Jan 2012

THE q-WAKIMOTO REALIZATION OF THE SUPERALGEBRAS $U_q(\widehat{sl}(N|1))$ AND $U_{q,p}(\widehat{sl}(N|1))$

February 1, 2012

TAKEO KOJIMA

Department of Mathematics and Physics, Graduate School of Science and Engineering,
Yamagata University, Jonan 4-3-16, Yonezawa 992-8510, Japan
kojima@yz.yamagata-u.ac.jp

Abstract

We give bosonizations of the superalgebras $U_q(\widehat{sl}(N|1))$ and $U_{q,p}(\widehat{sl}(N|1))$ for an arbitrary level $k \in \mathbf{C}$. We introduce the submodule by the ξ-η system, that we call the q-Wakimoto realization.

1 Introduction

Bosonizations are known to be a powerful method to construct correlation functions in not only conformal field theory [1], but also exactly solvable lattice models [2]. The quantum algebra $U_q(g)$ and the elliptic algebra $U_{q,p}(g)$ play an important role in exactly solvable lattice models. The level parameter k plays an important role in representation theory for $U_q(g)$ and $U_{q,p}(g)$. Bosonizations for an arbitrary level k are completely different from those of level $k=1$. In the case for level $k=1$, bosonizations have been constructed for quantum algebra $U_q(g)$ in many cases $g=(ADE)^{(r)}$, $(BC)^{(1)}$, $G_2^{(1)}$, $\widehat{sl}(M|N)$, $osp(2|2)^{(2)}$ [3, 6, 4, 5, 7, 8, 9, 10]. Using the dressing method developed in non-twisted algebra [11] and twisted algebra $A_2^{(2)}$ [12], we have bosonizations of the elliptic algebra $U_{q,p}(g)$ for $g=(ADE)^{(1)}, (BC)^{(1)}, G_2^{(1)}$ and $A_2^{(2)}$. In the case of an arbitrary level k, bosonizations have been constructed only for $U_q(\widehat{sl}(N))$ [14, 15, 16], $U_q(\widehat{sl}(2|1))$ [17], $U_{q,p}(\widehat{sl}(N))$ [11], and $U_{q,p}(\widehat{sl}(2|1))$ [19]. In this paper we give a bosonization of the quantum superalgebra $U_q(\widehat{sl}(N|1))$ for an arbitrary level k [20]. Using the dressing method developed in [19], we give a bosonization of the quantum superalgebra $U_{q,p}(\widehat{sl}(N|1))$ for an arbitrary level k. The level

k bosonizations on the boson Fock space of $U_q(\widehat{sl}(N))$ and $U_q(\widehat{sl}(N|1))$ [16, 17, 20] are not irreducible realizations. The construction of the irreducible highest weight module $V(\lambda)$ is nontrivial problem. We recall the non-quantum algebra $\widehat{sl}(2)$ case [21]. The irreducible highest weight module $V(\lambda)$ for the affine algebra $\widehat{sl}(2)$ was constructed from the Wakimoto realization on the boson Fock space [13] by the Felder complex. We recall the quantum algebra $U_q(\widehat{sl}(2))$ case [14, 15, 22]. The irreducible highest weight module $V(\lambda)$ for $U_q(\widehat{sl}(2))$ was constructed from the level k bosonizations on the boson Fock space [14, 15] by two steps; the first step is the resolution by the ξ-η system, and the second step is the resolution by the Felder complex [22, 21]. The submodule of the quantum algebra $U_q(\widehat{sl}(2))$, induced by the ξ-η system, plays the same role as the Wakimoto realization of the non-quantum algebra $\widehat{sl}(2)$. We would like to call this submodule induced by the ξ-η system "the q-Wakimoto realization". Constructions of the irreducible highest weight module $V(\lambda)$ for $U_q(\widehat{sl}(N))$ $(N \geq 3)$ and $U_q(\widehat{sl}(N|1))$ $(N \geq 2)$ are still an open problem. In this paper we study the ξ-η system and introduce the q-Wakimoto realization for the superalgebra $U_q(\widehat{sl}(N|1))$ and $U_{q,p}(\widehat{sl}(N|1))$.

This paper is organized as follows. In section 2, after preparing notations, we give the definition of the quantum superalgebra $U_q(\widehat{sl}(N|1))$ and the elliptic superalgebra $U_{q,p}(\widehat{sl}(N|1))$. In section 3 we give bosonizations of the superalgebras $U_q(\widehat{sl}(N|1))$ and $U_{q,p}(\widehat{sl}(N|1))$ for an arbitrary level k. In section 4 we introduce the q-Wakimoto realization of by the ξ-η system.

2 Superalgebra $U_q(\widehat{sl}(N|1))$ and $U_{q,p}(\widehat{sl}(N|1))$

In this section we recall the definitions of the quantum superalgebra $U_q(\widehat{sl}(N|1))$ [18] and the elliptic deformed superalgebra $U_{q,p}(\widehat{sl}(N|1))$ [19] for $N \geq 2$. We fix a complex number $q \neq 0, |q| < 1$. We set

$$[x, y] = xy - yx, \ \{x, y\} = xy + yx, \ [a]_q = \frac{q^a - q^{-a}}{q - q^{-1}}. \tag{2.1}$$

Let us fix complex numbers $r, k \in \mathbf{C}$, $\mathrm{Re}(r) > 0, \mathrm{Re}(r-k) > 0$. We use the abbreviation $r^* = r - k$. We set $p = q^{2r}$. We set the Jacobi theta functions

$$[u] = q^{\frac{u^2}{r} - u} \frac{\Theta_{q^{2r}}(q^{2u})}{(q^{2r}; q^{2r})_\infty^3}, \quad [u]^* = q^{\frac{u^2}{r^*} - u} \frac{\Theta_{q^{2r^*}}(q^{2u})}{(q^{2r^*}; q^{2r^*})_\infty^3}, \tag{2.2}$$

where we have used

$$\Theta_p(z) = (z; p)_\infty (pz^{-1}; p)_\infty (p; p)_\infty, \quad (z; p)_\infty = \prod_{m=0}^{\infty} (1 - p^m z). \tag{2.3}$$

The Cartan matrix $(A_{i,j})_{0 \leq i,j \leq N}$ of the affine Lie algebra $\widehat{sl}(N|1)$ is given by

$$A_{i,j} = (\nu_i + \nu_{i+1})\delta_{i,j} - \nu_i \delta_{i,j+1} - \nu_{i+1}\delta_{i+1,j}. \tag{2.4}$$

Here we set $\nu_1 = \cdots = \nu_N = +, \nu_{N+1} = \nu_0 -$.

2.1 Quantum superalgebra $U_q(\widehat{sl}(N|1))$

In this section we recall the definition of the quantum affine superalgebra $U_q(\widehat{sl}(N|1))$.

Definition 2.1 *[18] The Drinfeld generators of the quantum superalgebra $U_q(\widehat{sl}(N|1))$ are*

$$x_{i,m}^{\pm},\ h_{i,m},\ c,\quad (1\leq i\leq N, m\in \mathbf{Z}). \tag{2.5}$$

Defining relations are

$$c:\text{central},\ [h_i,h_{j,m}]=0, \tag{2.6}$$

$$[a_{i,m},h_{j,n}]=\frac{[A_{i,j}m]_q[cm]_q}{m}q^{-c|m|}\delta_{m+n,0}\quad (m,n\neq 0), \tag{2.7}$$

$$[h_i,x_j^{\pm}(z)]=\pm A_{i,j}x_j^{\pm}(z), \tag{2.8}$$

$$[h_{i,m},x_j^{+}(z)]=\frac{[A_{i,j}m]_q}{m}q^{-c|m|}z^m x_j^{+}(z)\quad (m\neq 0), \tag{2.9}$$

$$[h_{i,m},x_j^{-}(z)]=-\frac{[A_{i,j}m]_q}{m}z^m x_j^{-}(z)\quad (m\neq 0), \tag{2.10}$$

$$(z_1-q^{\pm A_{i,j}}z_2)x_i^{\pm}(z_1)x_j^{\pm}(z_2)=(q^{\pm A_{j,i}}z_1-z_2)x_j^{\pm}(z_2)x_i^{\pm}(z_1)\quad \text{for } |A_{i,j}|\neq 0, \tag{2.11}$$

$$x_i^{\pm}(z_1)x_j^{\pm}(z_2)=x_j^{\pm}(z_2)x_i^{\pm}(z_1)\quad \text{for } |A_{i,j}|=0, (i,j)\neq (N,N), \tag{2.12}$$

$$\{x_N^{\pm}(z_1),x_N^{\pm}(z_2)\}=0, \tag{2.13}$$

$$\begin{aligned}&[x_i^{+}(z_1),x_j^{-}(z_2)]=\frac{\delta_{i,j}}{(q-q^{-1})z_1z_2}\left(\delta(q^{-c}z_1/z_2)\psi_i^{+}(q^{\frac{c}{2}}z_2)-\delta(q^{c}z_1/z_2)\psi_i^{-}(q^{-\frac{c}{2}}z_2)\right),\\ &\quad \text{for}\quad (i,j)\neq (N,N),\end{aligned} \tag{2.14}$$

$$\{x_N^{+}(z_1),x_N^{-}(z_2)\}=\frac{1}{(q-q^{-1})z_1z_2}\left(\delta(q^{-c}z_1/z_2)\psi_N^{+}(q^{\frac{c}{2}}z_2)-\delta(q^{c}z_1/z_2)\psi_N^{-}(q^{-\frac{c}{2}}z_2)\right), \tag{2.15}$$

$$\begin{aligned}&\left(x_i^{\pm}(z_1)x_i^{\pm}(z_2)x_j^{\pm}(z)-(q+q^{-1})x_i^{\pm}(z_1)x_j^{\pm}(z)x_i^{\pm}(z_2)+x_j^{\pm}(z)x_i^{\pm}(z_1)x_i^{\pm}(z_2)\right)\\ &+(z_1\leftrightarrow z_2)=0\quad \text{for } |A_{i,j}|=1,\ i\neq N,\end{aligned} \tag{2.16}$$

where we have used $\delta(z)=\sum_{m\in\mathbf{Z}}z^m$. Here we have used the abbreviation $h_i=h_{i,0}$. We have set the generating function

$$x_j^{\pm}(z) = \sum_{m\in\mathbf{Z}}x_{j,m}^{\pm}z^{-m-1}, \tag{2.17}$$

$$\psi_i^{+}(q^{\frac{c}{2}}z) = q^{h_i}\exp\left((q-q^{-1})\sum_{m>0}h_{i,m}z^{-m}\right), \tag{2.18}$$

$$\psi_i^{-}(q^{-\frac{c}{2}}z) = q^{-h_i}\exp\left(-(q-q^{-1})\sum_{m>0}h_{i,-m}z^{m}\right). \tag{2.19}$$

2.2 Elliptic Superalgebra $U_{q,p}(\widehat{sl}(N|1))$

In this section we recall the definition of the elliptic superalgebra $U_{q,p}(\widehat{sl}(N|1))$.

Definition 2.2 *[19] The elliptic superalgebra $U_{q,p}(\widehat{sl}(N|1))$ is the associative algebra generated by the currents $E_j(z), F_j(z), H_j^{\pm}(z)$ $(1\leq j\leq N)$ and $B_{j,m}(1\leq j\leq N, m\in\mathbf{Z}_{\neq 0})$, h_j $(1\leq j\leq N)$ that satisfy the following relations.*

$$[h_i,B_{j,m}]=0,\ [B_{i,m},B_{j,n}]=\frac{[A_{i,j}m]_q[km]_q}{m}\frac{[r^*m]_q}{[rm]_q}\delta_{m+n,0}, \tag{2.20}$$

$$[h_i, E_j(z)] = A_{i,j} E_j(z), \ \ [h_i, F_j(z)] = -A_{i,j} F_j(z), \tag{2.21}$$

$$[B_{i,m}, E_j(z)] = \frac{[A_{i,j} m]_q}{m} z^m E_j(z), \tag{2.22}$$

$$[B_{i,m}, F_j(z)] = -\frac{[A_{i,j} m]_q}{m} \frac{[r^* m]_q}{[rm]_q} z^m F_j(z). \tag{2.23}$$

For $1 \leq i, j \leq N$ *such that* $(i,j) \neq (N,N)$ *they satisfy*

$$\left[u_1 - u_2 - \frac{A_{i,j}}{2}\right]^* E_i(z_1) E_j(z_2) = \left[u_1 - u_2 + \frac{A_{i,j}}{2}\right]^* E_j(z_2) E_i(z_1), \tag{2.24}$$

$$\left[u_1 - u_2 + \frac{A_{i,j}}{2}\right] F_i(z_1) F_j(z_2) = \left[u_1 - u_2 - \frac{A_{i,j}}{2}\right] F_j(z_2) F_i(z_1), \tag{2.25}$$

$$[E_i(z_1), F_j(z_2)] = \frac{\delta_{i,j}}{(q-q^{-1}) z_1 z_2} \left(\delta(q^{-k} z_1/z_2) H_i(q^r z_2) - \delta(q^k z_1/z_2) H_i(q^{-r} z_2) \right), \tag{2.26}$$

$$\{E_N(z_1), E_N(z_2)\} = 0, \ \ \{F_N(z_1), F_N(z_2)\} = 0, \tag{2.27}$$

$$\{E_N(z_1), F_N(z_2)\} = \frac{1}{(q-q^{-1}) z_1 z_2} \left(\delta(q^{-k} z_1/z_2) H_N(q^r z_2) - \delta(q^k z_1/z_2) H_N(q^{-r} z_2) \right). \tag{2.28}$$

For $1 \leq i, j \leq N$ *they satisfy*

$$H_i(z_1) H_j(z_2) = \frac{[u_2 - u_1 - \frac{A_{i,j}}{2}]^* [u_2 - u_1 + \frac{A_{i,j}}{2}]}{[u_2 - u_1 + \frac{A_{i,j}}{2}]^* [u_2 - u_1 - \frac{A_{i,j}}{2}]} H_j(z_2) H_i(z_1), \tag{2.29}$$

$$H_i(z_1) E_j(z_2) = \frac{[u_1 - u_2 + \frac{r^*}{2} + \frac{A_{i,j}}{2}]^*}{[u_1 - u_2 + \frac{r^*}{2} - \frac{A_{i,j}}{2}]^*} E_j(z_2) H_i(z_1), \tag{2.30}$$

$$H_i(z_1) F_j(z_2) = \frac{[u_1 - u_2 + \frac{r}{2} + \frac{A_{i,j}}{2}]}{[u_1 - u_2 + \frac{r}{2} - \frac{A_{i,j}}{2}]} F_j(z_2) H_i(z_1). \tag{2.31}$$

For $1 \leq i, j \leq N$,$(i \neq N)$ *such that* $|A_{i,j}| = 1$, *they satisfy the Serre relations.*

$$\begin{aligned}
&\left\{ E_i(z_1) E_i(z_2) E_j(z) \frac{\left(q^{2r^*+A_{i,j}} \frac{z}{z_1}; q^{2r^*}\right)_\infty \left(q^{2r^*+A_{i,j}} \frac{z}{z_2}; q^{2r^*}\right)_\infty}{\left(q^{2r^*-A_{i,j}} \frac{z}{z_1}; q^{2r^*}\right)_\infty \left(q^{2r^*-A_{i,j}} \frac{z}{z_2}; q^{2r^*}\right)_\infty} \left(\frac{z}{z_2}\right)^{\frac{1}{r^*} A_{i,j}} \right. \\
&-(q+q^{-1}) E_i(z_1) E_j(z) E_i(z_2) \frac{\left(q^{2r^*+A_{i,j}} \frac{z}{z_1}; q^{2r^*}\right)_\infty \left(q^{2r^*+A_{i,j}} \frac{z_2}{z}; q^{2r^*}\right)_\infty}{\left(q^{2r^*-A_{i,j}} \frac{z}{z_1}; q^{2r^*}\right)_\infty \left(q^{2r^*-A_{i,j}} \frac{z_2}{z}; q^{2r^*}\right)_\infty} \\
&\left. + E_j(z) E_i(z_1) E_i(z_2) \frac{\left(q^{2r^*+A_{i,j}} \frac{z_1}{z}; q^{2r^*}\right)_\infty \left(q^{2r^*+A_{i,j}} \frac{z_2}{z}; q^{2r^*}\right)_\infty}{\left(q^{2r^*-A_{i,j}} \frac{z_1}{z}; q^{2r^*}\right)_\infty \left(q^{2r^*-A_{i,j}} \frac{z_2}{z}; q^{2r^*}\right)_\infty} \left(\frac{z_1}{z}\right)^{\frac{1}{r^*} A_{i,j}} \right\} \\
&\times \frac{\left(q^{2r^*+A_{i,i}} \frac{z_2}{z_1}; q^{2r^*}\right)_\infty}{\left(q^{2r^*-A_{i,i}} \frac{z_2}{z_1}; q^{2r^*}\right)_\infty} z_1^{-\frac{1}{r^*}(A_{i,i}+A_{i,j})} + (z_1 \leftrightarrow z_2) = 0,
\end{aligned} \tag{2.32}$$

$$\begin{aligned}
&\left\{ F_i(z_1) F_i(z_2) F_j(z) \frac{\left(q^{2r-A_{i,j}} \frac{z}{z_1}; q^{2r}\right)_\infty \left(q^{2r-A_{i,j}} \frac{z}{z_2}; q^{2r}\right)_\infty}{\left(q^{2r+A_{i,j}} \frac{z}{z_1}; q^{2r}\right)_\infty \left(q^{2r+A_{i,j}} \frac{z}{z_2}; q^{2r}\right)_\infty} \left(\frac{z_2}{z}\right)^{\frac{1}{r} A_{i,j}} \right. \\
&-(q+q^{-1}) F_i(z_1) F_j(z) F_i(z_2) \frac{\left(q^{2r-A_{i,j}} \frac{z}{z_1}; q^{2r}\right)_\infty \left(q^{2r-A_{i,j}} \frac{z_2}{z}; q^{2r}\right)_\infty}{\left(q^{2r+A_{i,j}} \frac{z}{z_1}; q^{2r}\right)_\infty \left(q^{2r+A_{i,j}} \frac{z_2}{z}; q^{2r}\right)_\infty}
\end{aligned}$$

$$
+F_j(z)F_i(z_1)F_i(z_2)\frac{\left(q^{2r-A_{i,j}}\frac{z_1}{z};q^{2r}\right)_\infty\left(q^{2r-A_{i,j}}\frac{z_2}{z};q^{2r}\right)_\infty}{\left(q^{2r+A_{i,j}}\frac{z_1}{z};q^{2r}\right)_\infty\left(q^{2r+A_{i,j}}\frac{z_2}{z};q^{2r}\right)_\infty}\left(\frac{z}{z_1}\right)^{\frac{1}{r}A_{i,j}}\Bigg\}
$$

$$
\times\frac{\left(q^{2r-A_{i,i}}\frac{z_2}{z_1};q^{2r}\right)_\infty}{\left(q^{2r+A_{i,i}}\frac{z_2}{z_1};q^{2r}\right)_\infty}z_1^{\frac{1}{r}(A_{i,i}+A_{i,j})}+(z_1\leftrightarrow z_2)=0. \qquad (2.33)
$$

Here we have used $z_j = q^{2u_j}$.

3 Bosonization

In this section we give bosonizations of the superalgebras $U_q(\widehat{sl}(N|1))$ and $U_{q,p}(\widehat{sl}(N|1))$ for an arbitrary level k [17, 19, 20].

3.1 Boson

We fix the level $c = k \in \mathbf{C}$. We introduce the bosons and the zero-mode operators a_m^j, Q_a^j $(m \in \mathbf{Z}, 1 \leq j \leq N)$, $b_m^{i,j}, Q_b^{i,j}$ $(m \in \mathbf{Z}, 1 \leq i < j \leq N+1)$, $c_m^{i,j}, Q_c^{i,j}$ $(m \in \mathbf{Z}, 1 \leq i < j \leq N)$. The bosons $a_m^i, b_m^{i,j}, c_m^{i,j}$, $(m \in \mathbf{Z}_{\neq 0})$ and the zero-mode operators a_0^i, Q_a^i, $b_0^{i,j}, Q_b^{i,j}$, $c_0^{i,j}, Q_c^{i,j}$ satisfy

$$
[a_m^i, a_n^j] = \frac{[(k+N-1)m]_q[A_{i,j}m]_q}{m}\delta_{m+n,0}, \quad [a_0^i, Q_a^j] = (k+N-1)A_{i,j}, \qquad (3.1)
$$

$$
[b_m^{i,j}, b_n^{i',j'}] = -\nu_i\nu_j\frac{[m]_q^2}{m}\delta_{i,i'}\delta_{j,j'}\delta_{m+n,0}, \quad [b_0^{i,j}, Q_b^{i',j'}] = -\nu_i\nu_j\delta_{i,i'}\delta_{j,j'}, \qquad (3.2)
$$

$$
[c_m^{i,j}, c_n^{i',j'}] = \frac{[m]_q^2}{m}\delta_{i,i'}\delta_{j,j'}\delta_{m+n,0}, \quad [c_0^{i,j}, Q_c^{i',j'}] = \delta_{i,i'}\delta_{j,j'}. \qquad (3.3)
$$

We impose the cocycle condition on the zero-mode operator $Q_b^{i,j}$, $(1 \leq i < j \leq N+1)$ by

$$
[Q_b^{i,j}, Q_b^{i',j'}] = \delta_{j,N+1}\delta_{j',N+1}\pi\sqrt{-1} \qquad \text{for } (i,j) \neq (i',j'). \qquad (3.4)
$$

We have the following (anti) commutation relations

$$
\left[e^{Q_b^{i,j}}, e^{Q_b^{i',j'}}\right] = 0 \quad (1 \leq i < j \leq N, 1 \leq i' < j' \leq N), \qquad (3.5)
$$

$$
\left\{e^{Q_b^{i,N+1}}, e^{Q_b^{j,N+1}}\right\} = 0 \quad (1 \leq i \neq j \leq N). \qquad (3.6)
$$

We use the standard symbol of the normal orderings $::$. In what follows we use the abbreviations $b^{i,j}(z), c^{i,j}(z), b_\pm^{i,j}(z), a_\pm^j(z)$ given by

$$
b^{i,j}(z) = -\sum_{m\neq 0}\frac{b_m^{i,j}}{[m]_q}z^{-m} + Q_b^{i,j} + b_0^{i,j}\log z, \ c^{i,j}(z) = -\sum_{m\neq 0}\frac{c_m^{i,j}}{[m]_q}z^{-m} + Q_c^{i,j} + c_0^{i,j}\log z, \qquad (3.7)
$$

$$
b_\pm^{i,j}(z) = \pm(q-q^{-1})\sum_{\pm m>0}b_m^{i,j}z^{-m} \pm b_0^{i,j}\log q, \ a_\pm^j(z) = \pm(q-q^{-1})\sum_{\pm m>0}a_m^jz^{-m} \pm a_0^j\log q. \ (3.8)
$$

3.2 Quantum Superalgebra $U_q(\widehat{sl}(N|1))$

In this section we give a bosonization of the quantum superalgebra $U_q(\widehat{sl}(N|1))$ for an arbitrary level k.

Theorem 3.1 *[20] The Drinfeld currents $x_i^\pm(z)$, $\psi_i^\pm(z)$, $(1 \le i \le N)$ of $U_q(\widehat{sl}(N|1))$ for an arbitrary level k are realized by the bosonic operators as follows.*

$$\begin{aligned}
x_i^+(z) &= \frac{1}{(q-q^{-1})z}\sum_{j=1}^{i} :\exp\left((b+c)^{j,i}(q^{j-1}z)+\sum_{l=1}^{j-1}(b_+^{l,i+1}(q^{l-1}z)-b_+^{l,i}(q^{l}z))\right)\times \qquad (3.9)\\
&\times \left\{\exp\left(b_+^{j,i+1}(q^{j-1}z)-(b+c)^{j,i+1}(q^{j}z)\right)-\exp\left(b_-^{j,i+1}(q^{j-1}z)-(b+c)^{j,i+1}(q^{j-2}z)\right)\right\}:,\\
x_N^+(z) &= \sum_{j=1}^{N}:\exp\left((b+c)^{j,N}(q^{j-1}z)+b^{j,N+1}(q^{j-1}z)-\sum_{l=1}^{j-1}(b_+^{l,N+1}(q^{l}z)+b_+^{l,N}(q^{l}z))\right):, \qquad (3.10)
\end{aligned}$$

$$\begin{aligned}
x_i^-(z) &= q^{k+N-1}:\exp\left(a_+^i(q^{\frac{k+N-1}{2}}z)-b^{i,N+1}(q^{k+N-1}z)-b_+^{i+1,N+1}(q^{k+N-1}z)+b^{i+1,N+1}(q^{k+N}z)\right):\\
&+ \frac{1}{(q-q^{-1})z}\sum_{j=1}^{i-1}:\exp\left(a_-^i(q^{-\frac{k+N-1}{2}}z)\right)\times \qquad (3.11)\\
&\quad\times\exp\left((b+c)^{j,i+1}(q^{-k-j}z)+b_-^{i,n+1}(q^{-k-n}z)-b_-^{i+1,n+1}(q^{-k-n+1}z)\right)\\
&\quad\times\exp\left(\sum_{l=j+1}^{i}(b_-^{l,i+1}(q^{-k-l+1}z)-b_-^{l,i}(q^{-k-l}z))+\sum_{l=i+1}^{N}(b_-^{i,l}(q^{-k-l}z)-b_-^{i+1,l}(q^{-k-l+1}z))\right)\times\\
&\quad\times\left\{\exp\left(-b_-^{j,i}(q^{-k-j}z)-(b+c)^{j,i}(q^{-k-j+1}z)\right)-\exp\left(-b_+^{j,i}(q^{-k-j}z)-(b+c)^{j,i}(q^{-k-j-1}z)\right)\right\}:\\
&+ \frac{1}{(q-q^{-1})z}:\left\{\exp\left(a_-^i(q^{-\frac{k+N-1}{2}}z)+(b+c)^{i,i+1}(q^{-k-i}z)\right.\right.\\
&\quad\left.+\sum_{l=i+1}^{N}(b_-^{i,l}(q^{-k-l}z)-b_-^{i+1,l}(q^{-k-l+1}z))+b_-^{i,N+1}(q^{-k-N}z)-b_-^{i+1,N+1}(q^{-k-N+1}z)\right)\\
&- \exp\left(a_+^i(q^{\frac{k+N-1}{2}}z)+(b+c)^{i,i+1}(q^{k+i}z)\right.\\
&\quad\left.\left.+\sum_{l=i+1}^{N}(b_+^{i,l}(q^{k+l}z)-b_+^{i+1,l}(q^{k+l-1}z))+b_+^{i,N+1}(q^{k+N}z)-b_+^{i+1,N+1}(q^{k+N-1}z)\right)\right\}:\\
&- \frac{1}{(q-q^{-1})z}\sum_{j=i+1}^{N-1}:\exp\left(a_+^i(q^{\frac{k+N-1}{2}}z)\right)\times\\
&\quad\times\exp\left((b+c)^{i,j+1}(q^{k+j}z)+b_+^{i,N+1}(q^{k+N}z)-b_+^{i+1,N+1}(q^{k+N-1}z)\right)\times\\
&\quad\times\exp\left(\sum_{l=j+1}^{N}(b_+^{i,l}(q^{k+l}z)-b_+^{i+1,l}(q^{k+l-1}z))\right)\times\left\{\exp\left(b_+^{i+1,j+1}(q^{k+j}z)-(b+c)^{i+1,j+1}(q^{k+j+1}z)\right)\right.\\
&\quad\left.-\exp\left(b_-^{i+1,j+1}(q^{k+j}z)-(b+c)^{i+1,j+1}(q^{k+j-1}z)\right)\right\}:.
\end{aligned}$$

$$\begin{aligned}
x_N^-(z) &= \frac{1}{(q-q^{-1})z}\left\{\sum_{j=1}^{N-1}q^{j-1}:\exp\left(a_-^N(q^{-\frac{k+N-1}{2}}z)\right)\times\right. \qquad (3.12)\\
&\quad\times\exp\left(-b_+^{j,N+1}(q^{-k-j}z)-b^{j,N+1}(q^{-k-j-1}z)-\sum_{l=j+1}^{N-1}(b_-^{l,N}(q^{-k-l}z)+b_-^{l,N+1}(q^{-k-l}z))\right)\times\\
&\quad\times\left\{\exp\left(-b_+^{j,N}(q^{-k-j}z)-(b+c)^{j,N}(q^{-k-j-1}z)\right)-\exp\left(-b_-^{j,N}(q^{-k-j}z)-(b+c)^{j,N}(q^{-k-j+1}z)\right)\right\}:
\end{aligned}$$

$$
+ \quad q^{N-1} : \left\{ \exp\left(a_+^N(q^{\frac{k+N-1}{2}}z) - b^{N,N+1}(q^{k+N-1}z) \right) - \exp\left(a_-^N(q^{-\frac{k+N-1}{2}}z) - b^{N,N+1}(q^{-k-N+1}z) \right) \right\} : \Big\} .
$$

$$
\begin{aligned}
\psi_i^{\pm}(q^{\pm\frac{k}{2}}z) &= \exp\left(a_{\pm}^i(q^{\pm\frac{k+N-1}{2}}z) + \sum_{l=1}^{i} (b_{\pm}^{l,i+1}(q^{\pm(l+k-1)}z) - b_{\pm}^{l,i}(q^{\pm(l+k)}z) \right) && (3.13)\\
&\times \exp\left(\sum_{l=i+1}^{N} (b_{\pm}^{i,l}(q^{\pm(k+l)}z) - b_{\pm}^{i-1,l}(q^{\pm(k+l-1)}z) + b_{\pm}^{i,N+1}(q^{\pm(k+N)}z) - b_{\pm}^{i+1,N+1}(q^{\pm(k+N-1)}z) \right),\\
\psi_N^{\pm}(q^{\pm\frac{k}{2}}z) &= \exp\left(a_{\pm}^N(q^{\pm\frac{k+N-1}{2}}z) - \sum_{l=1}^{N-1} (b_{\pm}^{l,N}(q^{\pm(k+l)}z) + b_{\pm}^{l,N+1}(q^{\pm(k+l)}z)) \right). && (3.14)
\end{aligned}
$$

3.3 Elliptic Superalgebra $U_{q,p}(\widehat{sl}(N|1))$

In this section we give a bosonization of the elliptic superalgebra $U_{q,p}(\widehat{sl}(N|1))$ for an arbitrary level k, using the dressing deformation [20]. Let us introduce the zero-mode operators P_i, Q_i, $(1 \leq i \leq N)$ by

$$
[P_i, Q_j] = -\frac{A_{i,j}}{2} \quad (1 \leq i, j \leq N), \tag{3.15}
$$

where $(A_{i,j})_{1 \leq i,j \leq N}$ is the Cartan matrix of the classical $sl(N|1)$. In [20] the bosonization of the Drinfeld generator $h_{i,m}$ $(1 \leq i \leq N, m \in \mathbf{Z})$ is given by

$$
\begin{aligned}
h_{i,m} &= q^{-\frac{k+N-1}{2}|m|}a_m^i + \sum_{l=1}^{i}(q^{-(k+l-1)|m|}b_m^{l,i+1} - q^{-(k+l)|m|}b_m^{l,i}) && (3.16)\\
&+ \sum_{l=i+1}^{N}(q^{-(k+l)|m|}b_m^{i,l} - q^{-(k+l-1)|m|}b_m^{i+1,l}) + q^{-(k+N)|m|}b_m^{i,N+1} - q^{-(k+N-1)|m|}b_m^{i+1,N+1},\\
h_{N,m} &= q^{-\frac{k+N-1}{2}|m|}a_m^N - \sum_{l=1}^{N-1}(q^{-(k+l)|m|}b_m^{l,N} + q^{-(k+l)|m|}b_m^{l,N+1}). && (3.17)
\end{aligned}
$$

Let us set the boson $B_{j,m}$ $(1 \leq j \leq N, m \in \mathbf{Z}_{\neq 0})$ by

$$
B_{j,m} = \left\{ \begin{array}{ll} \frac{[r^*m]_q}{[rm]_q} h_{j,m} & (m > 0), \\ q^{k|m|}h_{j,m} & (m < 0). \end{array} \right. \tag{3.18}
$$

Theorem 3.2 *[19, 20] The currents $E_j(z), F_j(z), H_j^{\pm}(z)$ $(1 \leq j \leq N)$ of the elliptic superalgebra $U_{q,p}(\widehat{sl}(N|1))$ for an arbitrary level k are realized by the bosonic operators as follows.*

$$
E_j(z) = U_j^+(z)x_j^+(z)e^{2Q_j}z^{-\frac{1}{r^*}P_j}, \tag{3.19}
$$

$$
F_j(z) = x_j^-(z)U_j^-(z)z^{\frac{1}{r}(P_j+h_j)}, \tag{3.20}
$$

$$
H_j^{\pm}(z) = H_j(q^{\pm(r-\frac{k}{2})}z), \tag{3.21}
$$

$$
H_j(z) =: \exp\left(-\sum_{m \neq 0} \frac{B_{j,m}}{[r^*m]_q} z^{-m} \right) : e^{2Q_j}z^{-\frac{k}{rr^*}P_j + \frac{1}{r}h_j}. \tag{3.22}
$$

Here we have used the dressing operators $U_j^+(z), U_j^-(z)$ $(1 \leq j \leq N)$ given by

$$
U_j^+(z) = \exp\left(\sum_{m>0} \frac{q^{rm}}{[rm]_q} B_{j,-m}z^m \right), \ U_j^-(z) = \exp\left(-\sum_{m>0} \frac{q^{r^*m}}{[rm]_q} B_{j,m}z^{-m} \right). \tag{3.23}
$$

4 q-Wakimoto Realization

In this section we introduce the q-Wakimoto realization by the ξ-η system. We introduce the vacuum state $|0\rangle$ of the boson Fock space by

$$a_m^i|0\rangle=b_m^{i,j}|0\rangle=c_m^{i,j}|0\rangle=0 \quad (m\geq 0). \tag{4.1}$$

For complex numbers $p_a^i\in\mathbf{C}$ $(1\leq i\leq N)$, $p_b^{i,j}\in\mathbf{C}$ $(1\leq i<j\leq N+1)$, $p_c^{i,j}\in\mathbf{C}$ $(1\leq i<j\leq N)$, we set

$$\begin{aligned}&|p_a,p_b,p_c\rangle \\ =\ &\exp\left(\sum_{i,j=1}^N\frac{\mathrm{Min}(i,j)(N-1-\mathrm{Max}(i,j))}{(N-1)(k+N-1)}p_a^iQ_a^j-\sum_{1\leq i<j\leq N+1}p_b^{i,j}Q_b^{i,j}+\sum_{1\leq i<j\leq N}p_c^{i,j}Q_c^{i,j}\right)|0\rangle.\end{aligned} \tag{4.2}$$

It satisfies

$$a_0^i|p_a,p_b,p_c\rangle=p_a^i|p_a,p_b,p_c\rangle,\ b_0^{i,j}|p_a,p_b,p_c\rangle=p_b^{i,j}|p_a,p_b,p_c\rangle,\ c_0^{i,j}|p_a,p_b,p_c\rangle=p_c^{i,j}|p_a,p_b,p_c\rangle. \tag{4.3}$$

The boson Fock space $F(p_a,p_b,p_c)$ is generated by the bosons $a_m^i,b_m^{i,j},c_m^{i,j}$ on the vector $|p_a,p_b,p_c\rangle$. We set $U_q(\widehat{sl}(N|1))$-module $F(p_a)$ by

$$F(p_a)=\bigoplus_{\substack{p_b^{i,j}=-p_c^{i,j}\in\mathbf{Z}\ (1\leq i<j\leq N)\\ p_b^{i,N+1}\in\mathbf{Z}\ (1\leq i\leq N)}}F(p_a,p_b,p_c). \tag{4.4}$$

We have imposed the restriction $p_b^{i,j}=-p_c^{i,j}\in\mathbf{Z}$, because the $x_{i,m}^{\pm}$ change $Q_b^{i,j}+Q_c^{i,j}$. The module $F(p_a)$ is not irreducible representation. For instance, the irreducible highest weight module $V(\lambda)$ for $U_q(\widehat{sl}(2))$ was constructed from the similar space as $F(p_a)$ by two steps; the first step is the construction of the q-Wakimoto realization by the ξ-η system, and the second step is the resolution by the Felder complex [22]. In this paper we study the ξ-η system and introduce the q-Wakimoto realization for $U_q(\widehat{sl}(N|1))$. For $1\leq i<j\leq N$ we introduce

$$\eta^{i,j}(z)=\sum_{m\in\mathbf{Z}}\eta_m^{i,j}z^{-m-1}=:e^{c^{i,j}(z)}:,\quad \xi^{i,j}(z)=\sum_{m\in\mathbf{Z}}\xi_m^{i,j}z^{-m}=:e^{-c^{i,j}(z)}:. \tag{4.5}$$

The Fourier components $\eta_m^{i,j}=\oint\frac{dz}{2\pi\sqrt{-1}}z^m\eta^{i,j}(z)$, $\xi_m^{i,j}=\oint\frac{dz}{2\pi\sqrt{-1}}z^{m-1}\xi^{i,j}(z)$ $(m\in\mathbf{Z})$ are well defined on the space $F(p_a)$. They satisfy

$$\{\eta_m^{i,j},\xi_n^{i,j}\}=\delta_{m+n,0},\ \{\eta_m^{i,j},\eta_n^{i,j}\}=\{\xi_m^{i,j},\xi_n^{i,j}\}=0\quad(1\leq i<j\leq N), \tag{4.6}$$

$$[\eta_m^{i,j},\xi_n^{i',j'}]=[\eta_m^{i,j},\eta_n^{i',j'}]=[\xi_m^{i,j},\xi_n^{i',j'}]=0\quad(i,j)\neq(i',j'). \tag{4.7}$$

We focus our attention on the operators $\eta_0^{i,j},\xi_0^{i,j}$ satisfying $(\eta_0^{i,j})^2=0$, $(\xi_0^{i,j})^2=0$ and $\mathrm{Im}(\eta_0^{i,j})=\mathrm{Ker}(\eta_0^{i,j})$, $\mathrm{Im}(\xi_0^{i,j})=\mathrm{Ker}(\xi_0^{i,j})$. The products $\eta_0^{i,j}\xi_0^{i,j}$ and $\xi_0^{i,j}\eta_0^{i,j}$ are the projection operators

$$\eta_0^{i,j}\xi_0^{i,j}+\xi_0^{i,j}\eta_0^{i,j}=1. \tag{4.8}$$

We have a direct sum decomposition.

$$F(p_a) = \eta_0^{i,j}\xi_0^{i,j}F(p_a) \oplus \xi_0^{i,j}\eta_0^{i,j}F(p_a), \tag{4.9}$$

$$\mathrm{Ker}(\eta_0^{i,j}) = \eta_0^{i,j}\xi_0^{i,j}F(p_a),\ \mathrm{Coker}(\eta_0^{i,j}) = \xi_0^{i,j}\eta_0^{i,j}F(p_a). \tag{4.10}$$

Definition 4.1 *We introduce the subspace $\mathcal{F}(p_a)$ that we call the q-Wakimoto realization.*

$$\mathcal{F}(p_a) = \left(\prod_{1\leq i<j\leq N} \eta_0^{i,j}\xi_0^{i,j}\right) F(p_a) = \bigcap_{1\leq i<j\leq N} \mathrm{Ker}(\eta_0^{i,j}), \tag{4.11}$$

The dressing operators $U_i^{\pm}(z)$ and the zero-mode operators P_i, Q_i commute with $\eta_0^{i',j'}$. The bosonizations commute with the operators $\eta_0^{i',j'}$,$\xi_0^{i',j'}$ up to sign $\pm$.

Proposition 4.2 *The subspace $\mathcal{F}(p_a)$ is both $U_q(\widehat{sl}(N|1))$ and $U_{q,p}(\widehat{sl}(N|1))$ module.*

Let $\bar{\alpha}_i$, $\bar{\Lambda}_i$, $(1 \leq i \leq N)$ and $(\cdot|\cdot)$ be the simple roots, the fundamental weights, and the symmetric bilinear norm ; $(\bar{\alpha}_i, \bar{\alpha}_j) = A_{i,j}$, $(\bar{\alpha}_i, \bar{\Lambda}_j) = \delta_{i,j}$. It is expected that we have the irreducible highest weight module $V(\lambda)$ with the highest weight λ, whose classical part $\bar{\lambda} = \sum_{j=1}^N p_a^i \bar{\Lambda}_i$, by the Felder complex of the q-Wakimoto realization. We would like to report this problem for $U_q(\widehat{sl}(N))$ and $U_q(\widehat{sl}(N|1))$ in the future publication.

Acknowledgements

This work is supported by the Grant-in-Aid for Scientific Research **C** (21540228) from Japan Society for Promotion of Science. The author would like to thank the organizing committee of the 9-th International Workshop "Lie Theory and its application in Physics" for an invitation to Bulgaria.

References

[1] P.Bouwknegt, J.McCarthy and K.Pilch, *Prog.Theor.Phys.***102** (1990) 67-135.

[2] M.Jimbo and T.Miwa, *Algebraic Analysis of Solvable Lattice Models*, CBMS Regional Conference Series in Mathematics **85** (American Mathematical Society), 1994.

[3] I.B.Frenkel and N.Jing, *Proc.Natl.Acad.Sci.* **85** (1988) 9373-9377.

[4] D.Bernard, *Lett.Math.Phys.***17** (1989) 239-245.

[5] N.Jing, Y.Koyama and K.Misra, *Selecta Math.***5** no. 2 (1999) 243-255.

[6] N.Jing, *Invent.Math.***102** (1990) 663-690.

[7] N.Jing, *Proc.Amer.Math.Soc.***127** no.1 (1999) 21-27.

[8] K.Kimura, J.Shiraishi and J.Uchiyama, *Comm. Math. Phys.* **188** no. 2 (1997) 367-378.

[9] Y.-Z.Zhang, *J.Math.Phys.***40** no.11 (1999) 6110-6124.

[10] W.-L.Yang and Y.-Z.Zhang, *Phys.Lett.***A261** (1999) 252-258.

[11] M.Jimbo, H.Konno, S.Odake and J.Shiraishi, *Commun.Math.Phys.* **199** (1999) 605-647.

[12] T.Kojima and H.Konno, *J.Math.Phys.* **45** (2004) 3146-3179.

[13] M.Wakimoto, *Commun.Math.Phys.***104** no.4 (1986) 605-609.

[14] A.Matsuo, *Commun.Math.Phys.***160** (1994) 33-48.

[15] J.Shiraishi, *Phys.Lett.***A171** (1992) 243-248.

[16] H.Awata, S.Odake and J.Shiraishi, *Commun.Math.Phys.* **162** no. 1 (1994) 61-83.

[17] H.Awata, S.Odake and J.Shiraishi, *Lett. Math. Phys.***42** no. 3 (1997) 271-279.

[18] H.Yamane, *Publ.Res.Inst.Math.Sci.* **35** (1999) 321-390.

[19] T.Kojima, *J.Phys.***A44**:Math.Theor. (2011) 485205(23pp).

[20] T.Kojima, *J.Math.Phys.***53** (2012) 013515(1-15).

[21] D.Bernard and G.Felder, *Commun.Math.Phys.***127** (1990) 145-168.

[22] H.Konno, *Mod.Phys.Lett.***A9** (1994) 1253-1265.

[23] Y.-Z.Zhang and M.D.Gould, *J.Math.Phys.***41** (2000) 5577-5291.